PAV 1800

Refrigerated *Pavlova*

For Bivalve Larvae, Spat, Broodstock, and more



PRODUCT DESCRIPTION:

Pav 1800 is a refrigerated, whole-cell concentrate of *Pavlova* sp. microalgae. Pav 1800 is very high in EPA, DHA and carotenoids (Strain: Pavlova sp, CCPM 459).

APPLICATIONS:

Effective for all life stages: **first feeding larvae**, **setting**, **spat** and **broodstock conditioning**. Produces increased growth and survival rates. Common applications:

- Peak Load Supplement
- 100% live algae replacement
- Remote set
- Back-up feed supply
- For use with open tanks, silos, bottle upwellers, header tanks, cultch tanks and depuration tanks
- Effective for many species including artemia, crab, sea cucumber and some copepods

BENEFITS:

- Biosecure
- Exceptionally Clean
- Easy to use
- Always available
- High levels of balanced lipids (DHA, EPA), protein, and carotenoids
- Has been Indicated to sychronize setting
- Effective feedd supplement and live algae replacement

APPEARANCE, PACKAGING & STORAGE:

- Viscous brown liquid concentrate
- 1-quart bottle, 10-liter Cubitainer®
- 4-month shelf life when refrigerated @ 0-5°C
- Do not freeze



Reed Mariculture
ENSURING HATCHERY SUCCESS

Producers of the Instant Algae,® Reef Nutrition® and APBreed® product lines •
Distributors of Otohime® and Mizuho® products for Marubeni Nisshin, Japan • Distributors of fresh Chlorella for Chlorella Ltd, Japan

Instant Algae®— Algae When You Need It®

TECHNICAL DATA:

Physical Properties

Algal Cell Size..... 4–7 microns
Cell Density..... ~3.9 Billion cells per ml
Algal Biomass..... 80g Dry Biomass/L

Composition of Dry Algal Biomass (Typical)

Protein.....51%
Lipids.....19%
DHA..... 6.0% (% of lipids)
EPA.....20.7% (% of lipids)
ArA..... 0.6% (% of lipids)
Carbohydrate.....22.0%
Ash.....8.0%

Composition of Liquid Product

Protein..... > 4.0%
Lipids..... > 1.5%
Carbohydrate..... > 1.7%
Ash..... < 8.0%
Moisture..... < 87%

Microbial Specifications

Coliform Bacteria..... <0.3 mpn/ml
Salmonella..... Negative
Known pathogenic marine bacteria... Negative

DIRECTIONS:

1. When water is circulated with a centrifugal pump, such as with an upweller, add in front of the pump intake and the pump will disperse the algae cells.
2. When feeding into open tanks or header tanks, pre-dilute the algae 10:1 with system water.
   a. Add system water to a small bucket.
   b. Add Pav 1800 and stir.
   c. DO NOT BLEND; it is unnecessary and may damage the algae cells.
   d. Pour into larvae or spat tank.
   e. Pouring through a 20–micron screen can improve dispersion.
3. Feed according to the table. Adjust dosage to meet the needs of your hatchery and larvae.
4. Refrigerate bottle when not in use.

TYPICAL FEEDING RATION

Larvae:

Feed according to shell length. Shell length on a given day is an estimate. Larviculture may run up to 20 days.

Daily Feed Rates per million larvae, *Crassostrea virginica* (from FAO Protocol 'Hatchery culture of bivalves 2004')

Shell Length	Day	PAV 1800 (ml)	=	Live *Isochrysis*(L) @4x10^6
75	2	0.4	=	0.4
95	3	0.5	=	0.6
100	4	0.6	=	0.7
115	5	2.0	=	2.1
130	6	3.4	=	3.6
145	7	4.8	=	5.1
160	8	6.1	=	6.6
190	9	8.8	=	9.5
220	10	11.6	=	12.4
240	11	13.4	=	14.4
260	12	15.4	=	16.3
270	13	16.2	=	17.3
280	14	17.1	=	18.3

Spat:

Feed by Weight

• 0.6ml Pav 1800 per gram live weight of spat per day
OR. *See website for 'Feed by size' table*

Broodstock Conditioning and Fattening:

Typical Broodstock Feed Rates

• 0.03–0.05 ml Pav 1800 per gram wet meat weight per day*
OR:
• 0.3–1.1 ml Shellfish Diet per adult animal per day (depending on feed ration and size*)
** Provides between 2% and 4% dry weight feed for dry weight of meat*

For details on feeding and batch or automated dispensing, visit: InstantAlgaeShellfish.com

Reed Mariculture
ENSURING HATCHERY SUCCESS

Producers of the Instant Algae,® Reef Nutrition® and APBreed® product lines •
Distributors of Otohime® and Mizuho® products for Marubeni Nisshin, Japan • Distributors of fresh Chlorella for Chlorella Ltd, Japan